

AGB超新星爆発における元素合成

和南城, 野本 (IPMU)

H. -Th. Janka, F. S. Kitaura, B. Müller (MPA)

arXiv:0810.3999

supernovae from 8-10 M_{\odot} stars

AGB SN from of an ONeMg core (up to ~30% of all SNe)

Nomoto 1984, 1987; Miyaji & Nomoto 1987

⇒ progenitor of Crab SN?

Nomoto et al. 1982; Davidson et al 1982

→ carbon-rich?? >9.5 M_{\odot} ??

MacAlpine & Satterfield 2008

⇒ low-luminosity SNe?

SN1997D, Chugai & Utrobin 2000

SN2003gd, Hendry et al. 2005

→ < 17 M_{\odot} for all SNeIIP??

Smartt et al. 2008


→ new SNe class of
SN2008S-like transients
with AGB progenitors??

Prieto et al. 2008; Thompson et al. 2008




Crab Nebula, hubblesite.org

origin of r-process nuclei?



(Ishimaru, Wanajo, Aoki, & Ryan 2004)



➡ chemical evolution studies favor the low-mass end ($8-10M_{\odot}$ stars) Ishimaru & Wanajo 1999; Ishimaru et al. 2004

➡ r-process models related to $8-10M_{\odot}$ stars Wanajo et al. 2003; Ning et al. 2007 but Janka et al. 2008; Hoffman et al. 2008

self-consistent explosion of a $9M_{\odot}$ star!

weak explosion by neutrino heating

Kitaura, Janka, & Hillebrandt 2006

→ the only case (1D)
with accurate ν transport

→ $E_{\text{exp}} = 1 \times 10^{50}$ erg


$M_{\text{ej}} = 0.014 M_{\odot}$

$s = 10\text{-}30 \text{ k/m}_u$


$Y_e = 0.46\text{-}0.53$

Nucleosynthesis
in the SN with
no parameters

→ most reliable SN yields
to date!?



initial compositions (Y_e)





nucleosynthesis

⇒ Fe (as ^{56}Ni) production
at $Y_e \geq 0.49$
($0.002-0.004M_\odot$)

⇒ ^{64}Zn and light p-nuclei
(e.g., ^{92}Mo) at $Y_e = 0.46-0.49$
Hoffman et al. 1996; Wanajo 2006


⇒ marginal vp-process
at $Y_e > 0.50$
Flohrich et al. 2006;
Pruet et al. 2006
Wanajo 2006

⇒ no r-process (up to ^{90}Zr)
at least $t_{\text{pb}} < 1\text{s}$ with 1D
Hoffman et al. 2008



overproduction factors



➔ strong overproduction of ^{90}Zr at $Y_e \sim 0.47$



mass-averaged yields

→ strong overproduction of ^{90}Zr , high Ni/Fe (~ 10 solar)

→ little production of ^{56}Ni ($=0.002-0.004M_{\odot}$ in mass fraction)



contribution to the Galaxy

$$\frac{f}{1-f} = \frac{X(^{90}\text{Zr})_{\square} / X(^{16}\text{O})_{\square}}{M(^{90}\text{Zr}) / M(^{16}\text{O})_{\text{other}}} = 0.029$$

f : fraction of ONeMg SNe relative to all SNeII/Ibc

$M(^{16}\text{O})_{\text{other}} = 1.5M_{\square}$: average ejecta mass of ^{16}O


per event from SNe ($> 10M_{\square}$)

⇒ $f = 0.028$

⇒ 81% of solar ^{90}Zr is from s-process

⇒ no more than 1% of all core-collapse supernovae....
in agreement with Hoffman et al. 2008


initial compositions (Y_e)



mass-averaged yields

➔ only 2% increase of $Y_{e,\min}$ cures the overproduction of ^{90}Zr !!

➔ robust production of ^{64}Zn and p-nuclei ^{74}Se , ^{78}Kr , ^{84}Sr , ^{92}Mo



contribution to the Galaxy

$$\frac{f}{1-f} = \frac{X(^{64}\text{Zn})_{\square} / X(^{16}\text{O})_{\square}}{M(^{64}\text{Zn}) / M(^{16}\text{O})_{\text{other}}} = 0.39$$

f : fraction of ONeMg SNe relative to all SNeII/Ibc

$M(^{16}\text{O})_{\text{other}} = 1.5M_{\square}$: average ejecta mass of ^{16}O


per event from SNe ($> 10M_{\square}$)

⇒ $f = 0.28$


⇒ origin solar ^{64}Zn (dominant isotope) is uncertain

⇒ up to ~ 30% of all core-collapse supernovae !!!
(consistent with 8-10 M_{\odot})

2D effect ? (preliminary)



(Kitaura et al. 2006)



(Muller & Janka in prep.)

conclusions

- ⇒ up to ~30% (~8-10 M_{\odot}) of all core-collapse supernovae possible origin of ^{64}Zn and p-nuclei ^{74}Se , ^{78}Kr , ^{84}Sr , ^{92}Mo
- ⇒ little ^{56}Ni production (=0.002-0.004 M_{\odot} in mass fraction) consistent with the Crab SN (with high Ni/Fe ~10 solar), low-luminosity SNeIIP (SN1997D, SN2003gd,), new class of luminous transients (SN2008S,)
- ⇒ carbon-rich gas of Crab, AGB progenitor of SN2008S mass range ~9.5-10 M_{\odot} (~7-8% of all SNe)